

## Effectiveness of Cardiac Rehabilitation on Co-Morbid Psychiatric Symptoms of Ischemic Heart Patients

Saeed Shoja Shafti MD<sup>1\*</sup> and Masuod Ahmadie MD<sup>2</sup>

<sup>1</sup>Professor of Psychiatry, University of Social Welfare and Rehabilitation Sciences (USWR), Razi Psychiatric Hospital, Tehran – Iran

<sup>2</sup>Cardiologist, Gilan – Iran

**\*Corresponding Author:** Saeed Shoja Shafti MD, Professor of Psychiatry, University of Social Welfare and Rehabilitation Sciences (USWR), Razi Psychiatric Hospital, Tehran – Iran.

**Received:** February 06, 2019; **Published:** February 15, 2019

### Abstract

**Introduction:** The high prevalence of co-morbidity among cardiac patients may result in undesired consequences, which certainly can compromise their managements. Cardiac rehabilitation can be an appropriate adjunctive intervention for amelioration of such morbid processes. In the current study we evaluated the effectiveness of cardiac rehabilitation on amelioration of depressive symptoms in a group of ischemic heart patients with co-morbid depressive illness.

**Methods and Materials:** An open comparative study had been performed between a group of ischemic cardiac patients with depressive symptoms, who received cardiac rehabilitation programs, in addition to their standard treatments (n=30), in comparison with a similar group of ischemic patients who had received only the customary managements (n=30).

**Result:** Significantly, more patients in the target group (56.6%) gained benefit from cardiac rehabilitation, in comparison with only 30 % of them in the control group (P<0.05), which was evident especially with respect to symptoms like anxiety, hopelessness, helplessness, hypochondriasis, depressed mood and work activities.

**Conclusion:** Consistent with encouraging outcome of present study, cardiac rehabilitation can be recommended as a supplementary technique, in addition to other standard treatments, in management of ischemic heart patients with co-morbid depressive illness.

Volume 3 Issue 1 February 2019

© All Copy Rights are Reserved by Saeed Shoja Shafti MD and Masuod Ahmadie MD.

### Introduction

Cardiac rehabilitation, as one of the known tactics for functional restoration of cardiac functions, according to some studies may have important healing or preventive effects with respect to psychiatric problems, like depression or anxiety, in patients who have suffered cardiovascular accidents. Psychiatric problems can exacerbate cardiovascular dysfunction a number of mechanisms [1,2]. For example, in depressive patients, the somatization of symptoms can reinforce the somatic complaints of ischemic heart patients or, alternately, by reducing the secretion of beta-endorphin the chest pain and its severity can really be increased [2].

**Citation:** Saeed Shoja Shafti MD and Masuod Ahmadie MD. "Effectiveness of Cardiac Rehabilitation on Co-Morbid Psychiatric Symptoms of Ischemic Heart Patients". *Current Opinions in Neurological Science* 3.1 (2019): 637-643.

Non-compliance with diet and prescribed medications [3], disturbed relationship with caretaker, imposed unnecessary restriction in social and occupational activities due to cognitive distortions or psychomotor retardation, sleep problems, irritability or agitation with its dangerous homodynamic sequels, lowered threshold and vulnerability to cardiovascular accidents due to arrhythmias and decreased sympathetic and increased parasympathetic activity, stress induced coronary vasospasm and platelet aggregation due to increased epinephrine and decreased substance P, are among the major sequels of depression, which has a life-time prevalence of 26% among ischemic cardiac patients [1]. In a prevalence survey in Tehran, there was a high rate of psychiatric co-morbidity (64%) among 101 ischemic heart patients who had been admitted in the coronary care unit (CCU) of an academic general hospital [4].

All of those psychiatric problems had been started many months or years before admission in the CCU. 42.57% of them had co-morbid depression, 9.90% had an anxiety disorder, and 10.89% were suffering from Not Otherwise Specified (NOS) problems. According to the findings of that study, integration of psychiatric co-morbidity and biological risk factors (hypertension, diabetes mellitus, cigarette smoking, and hyperlipidemia) could increase the risk of cardiac events up to three folds [4]. In general, by considering the additional risk (16-20%) of occurrence of major depressive disorder after myocardial infarction (half of them show depressive symptoms 3 months after infarction) with its fivefold increase in mortality in the next six months, we can understand the importance of psychiatric co-morbidity in coronary artery disease.

Even hopelessness as a single symptom in depression, disregard to other symptoms, increase the risk of morbidity 1.5-2.5 fold [1]. Moreover, three community-based studies have shown a significant relationship between anxiety disorders and sudden cardiac death. One study of 1,457 male subjects followed up over 6 years found the risk of cardiac-related death elevated almost fourfold in those with phobic anxiety [5]. Acutely stressful events, as well, dramatically increase cardiac morbidity and mortality. In a prospective study of 95,647 Finnish men and women, the risk of all-cause mortality in the first week following spousal death was two times normal. The risk of any ischemic heart disease event in that period was elevated 2.3-fold in men and 3.5-fold in women, an effect independent of age [6].

Finally, regarding relationship between Type A behavior pattern (TABP)—time urgency, hostility, and achievement-striving-competitiveness—and the risk of coronary heart disease, while a 1987 meta-analysis documented the weakening of the association of type A behavior with coronary heart disease [7], the controversies still go on [8–10]. Therefore, cardiac rehabilitation with its supportive, cognitive and behavioral aspects seems to have important psychotherapeutic benefit in accompany with other standard treatments. In this research we evaluated the efficacy of cardiac rehabilitation on amelioration of depressive symptoms. In the current study we evaluated the effectiveness of cardiac rehabilitation on amelioration of depressive symptoms in a group of ischemic heart patients with co-morbid depressive illness.

## **Methods and Materials**

In an open trial, 30 randomly chosen patients, among referred ones to a cardiac rehabilitation center with the history of a recent myocardial infarction (in the last month), who were suffering from co-morbid depression, too (Table-1), had been delivered a rehabilitation program including:

1. Explanation of the nature of cardiovascular diseases,
2. Description of the essential process of management,
3. Recommendations for taking necessary steps for dealing with existing risk factors and stresses (stress management), and
4. Appropriate exercises according to the patient's compliance with accentuation on restoring the ambulatory status and functional activities as soon as possible after aforesaid cardiac event, in addition to their standard treatments.

All of these programs had been accomplished by the rehabilitation team, including: cardiologist (head of team), psychiatrist, psychologist, registered nurse and physiotherapist. The program had been accomplished on cardiac patients, out-patiently, within four weeks of being discharged from the hospital, and the session, as well, had been performed once per week (30-45 minutes), for duration of eight weeks per patient. The diagnosis of depression in both groups had been certified by a psychiatrist according to the DSM IV-TR

criteria and had been scored by Hamilton Rating Scale for Depression (HRS-D) [11] at the beginning of the program and then every two weeks during the trial.

Inclusion Criteria	Exclusion Criteria
Ischemic heart patients with myocardial infarction in the last month, plus co-morbid depression, which have been started no less than one month before cardiac event.	<ol style="list-style-type: none"> <li>1. Non-ischemic cardiac patients.</li> <li>2. Ischemic cardiac patients without depression.</li> <li>3. Ischemic cardiac patients without myocardial infarction.</li> <li>4. Infarction more than one month ago.</li> <li>5. Depression shorter than one month before infarction.</li> </ol>

**Table 1:** Inclusion and exclusion criteria for selecting patients.

All of these programs had been accomplished by the rehabilitation team, including: cardiologist (head of team), psychiatrist, psychologist, registered nurse and physiotherapist. The program had been accomplished on cardiac patients, out-patiently, within four weeks of being discharged from the hospital, and the session, as well, had been performed once per week (30-45 minutes), for duration of eight weeks per patient. The diagnosis of depression in both groups had been certified by a psychiatrist according to the DSM IV-TR criteria and had been scored by Hamilton Rating Scale for Depression (HRS-D) [11] at the beginning of the program and then every two weeks during the trial.

The HRS-D (HAM-D) is an observer-rated scale consisting of 17 to 21 items. Rating is made on the basis of the clinical interview, plus any additional available information by other resources such as staff or family members. The items are rated on either a 0 to 4 (0=none/absent and 4=most severe) or a 0 to 2 spectrum (0=absent/none and 2=severe). Depressed Mood, Feelings of Guilt, Suicide, Insomnia, activities, retardation, agitation, anxiety psychic, anxiety somatic, hypochondriasis, changes of weight, insight, diurnal variation, depersonalization and derealization, paranoid ideations, and obsessive or compulsive symptoms are the items which are scored in HRS-D.

The control group, too, included 30 randomly chosen ischemic cardiac patients with similar co-morbid psychopathology, among the referred ones, who received only the standard treatments (antihypertensive, anti-angina, and antidepressant medications). All of the aforesaid patients in both groups had been prescribed antidepressants from no less than one month before their entrance into the trial. Total duration of study was around eighteen months, from August of 2000 till February of 2002.

### Statistical Analysis

Patients were compared on baseline characteristics by means of t-tests. Treatment efficacy also was analyzed by 'Comparison of Proportions'. Statistical significance was defined as a 2-sided p value < or = to 0.05. MedCalc, version 9.4.1.0, was used as statistical software tool for analysis.

### Results

According to data, the age of the patients in the target group was between 34 and 73 (mean=51.46 ± 8.56) years old and all of them were male. Duration of their ischemic heart disease was between one month and 24 years (mean=11.35 ± 2.79 years). These parameters in the control group were 31 to 66 (mean=49.88 ± 9.31) years old and one month up to 19 years (mean=10.54 ± 4.16 years) respectively, and all of them were male, too. 'Comparison of means' did not show any significant difference with respect to age (t = 0.684, P < 0.49, CI = -3.04, 6.20) or duration of ischemic illness (t = 0.886, P < 0.37, CI = -1.03, 2.65) between two groups.

According to the scores of HRS-D their depression had been classified into three groups as follows: Mild (10-13), Mild to Moderate (14-17), and Moderate to Severe (>17) (Table - 2). 56.6 % (n=17) of the patients in the target group and 30 % (n=9) of them in the control group showed some improvement in some or most of their sub-scales. 'Comparison of Proportions' showed a significant difference with respect to response to the aforesaid rehabilitation program between two groups (z = 2.08, p-value <0.03, CI 95% = 0.015, 0.517).

The items which exhibited the most influence, in this regard, in the target group were: depressed mood (2 to 1 or 1 to 0), work and activities (1 to 0), helplessness (2 to 1 or 1 to 0), hopelessness (2 to 1 or 1 to 0), anxiety psychic (1 to 0), and hypochondriasis (2 to 1 or 1 to 0) in 20%, 20%, 23.33%, 26.66%, 36.67% and finally 23.33% of the patients, respectively. Psychic anxiety was the most influenced one in the target group (Table 3).

Outcome measure (HRS-D) Group	Mild (10-13)	Mild to Moderate (14-17)	Moderate to Severe (>17)	Total
Target	17 (56.67%)	8 (26.67%)	5 (16.67%)	30 (100%)
Control	14 (46.67%)	10 (33.33%)	6 (20%)	30 (100%)

Table 2: Classification of patients according to the severity of depression.

HRS-D ITEMS Patients	Depressed Mood (2 to 1-1 to 0)	Work Activities (1 to 0)	Helplessness (2 to 1-1 to 0)	Hopelessness (2 to 1-1 to 0)	Anxiety Psychic (1 to 0)	Hypochondriasis (2 to 1-1 to 0)
1	+	+	+	+	+	+
2	+	+	+			
3		+		+		+
4			+	+	+	
5			+	+	+	
6	+		+	+	+	
7		+				+
8		+				
9					+	
10	+			+	+	
11					+	+
12	+		+	+		
13					+	
14					+	+
15	+				+	+
16			+	+	+	
17		+				+
Total	6	6	7	8	11	7

Table 3: Characteristics of improved symptoms in the target group.

Most improvements had appeared in the last two months. No important improvement was evident in the remaining items. Besides, all of these positive responses had been palpable in the Mild and Mild to Moderate classes of that group. In the control group, too, there were some improvements in the depressed mood (16.66%), work activities (10%), helplessness (10%), hopelessness (10%), anxiety psychic (16.66%), hypochondriasis (10%), agitation (10%), and retardation (6.66%). The psychic anxiety and depressed mood also were the items with the greatest improvement in the control group (Table 4), which had appeared mostly in the last ten weeks of the trial among the first two classes of the patients (Figure 1).

HRS-D items Patients	Depressed mood (2 to 1-1 to 0)	Work Activities (1 to 0)	Helplessness (2 to 1-1 to 0)	Hopelessness (2 to 1-1 to 0)	Anxiety Psychic (1 to 0)	Hypochondriasis (2 to 1-1 to 0)	Agitation (1 to 0)	Retardation (1 to 0)
1	+	+	+	+	+		+	+
2	+				+		+	
3			+	+		+		
4	+	+			+			
5			+	+				
6						+	+	
7		+			+			+
8	+				+			
9	+					+		
Total	5	3	3	3	5	3	3	2

Table 4: Characteristics of improved symptoms in the control group.




Figure 1: Numerical differences re positive response according to HRS-D.

Discussion

Anxiety and depression are more prevalent in patients with cardiovascular disease than in the general population [12]. A study of 200 patients who had suffered a first myocardial infarction (MI) found a 1-year cumulative incidence of depression of 25% [13]. Also, in patients with congestive heart failure (CHF), three studies have found the prevalence of depression to be approximately 20% [14]. Although the epidemiology of anxiety disorders in CAD is not as well studied as that of depression, the incidence of anxiety symptoms in patients with acute coronary disease in cardiac care units is approximately 50% [15]. Besides, depression is linked to a large number of major risk factors for CAD or cardiac-related mortality, including cigarette smoking, diabetes, and obesity.

Prospective studies indicate that depression is an independent risk factor in the development of these behaviors or conditions and also strong independent risk factor for the development of cardiovascular disease, of acute coronary events, and of mortality from cardiac illness. In general, studies show that the relative risk of incident cardiac disease in healthy individuals with depression or symptoms of depression is about 1.5 to 2, depending on which cardiac endpoint is used [16,17]. Also, findings of a recent study on 2,111 incident myocardial infarctions, in a large Norwegian population based cohort, showed that self-reported symptoms of depression and anxiety, especially if recurrent, were moderately associated with the risk of acute myocardial infarction [18].

Citation: Saeed Shoja Shafti MD and Masuod Ahmadie MD. "Effectiveness of Cardiac Rehabilitation on Co-Morbid Psychiatric Symptoms of Ischemic Heart Patients". *Current Opinions in Neurological Science* 3.1 (2019): 637-643.

Though there was not any substantial wide-ranging reduction in HRS-D in the target group and, likewise, there was not any improvement with respect to severe symptoms, and over again all of the positive responses were slight (1 to 0 or 2 to 1), nevertheless the better response of depressive symptoms in the target group in comparison with the control one, which was evident significantly in 'comparison of proportions', had revealed an adjunctive remedial influence of cardiac rehabilitation on this group of cardiac patients; an outcome similar to the findings of Lavie, *et al.* [19], Milani, *et al.* [20], Linden, *et al.* [21] and Johnston, *et al.* [22]. But while methodology of most of the said scholars addressed mainly post-MI mood problems (in contrary to present study which had targeted premorbid mood difficulties just before occurrence of cardiac event), but according to the findings of the current try-out its assistance seems to be generalizable to co-morbid psychiatric symptoms regardless to its primary or secondary process.

Moreover, it should not be disregarded that many of the depressive symptoms which appear after such serious events are merely adjustment disorder, not a full-blown major depressive disorder. Therefore, while negative cognitive processing of cardiac patients can be strengthened by the influence of depressive mood and, consecutively, reinforcement of somatic problems and deterioration of their survival, cardiac rehabilitation may decline this negative process by its verbal (explanatory) and nonverbal (physiotherapeutic) components and increase patient's self-esteem and self – competency, which are antagonist of self-depreciation and pessimistic point of view [23].

Several studies have reported improvement in psychological stress in patients with coronary heart disease who have attended cardiac rehabilitation: one recent US observational study of a number of patients with heart failure (left ventricular ejection fraction <45%) reported a decrease in symptoms of depression by 40% after exercise training cardiac rehabilitation [20]. Also depressed patients who completed their cardiac rehabilitation had a 59% lower mortality compared with depressed dropout patients who did not undergo cardiac rehabilitation [20].

So, cardiac rehabilitation seems to be a fair adjuvant therapeutic approach which, from a psychotherapeutic standpoint, can decrease the negative cognitive distortion resulting from the depression, co-morbid with ischemic heart disease, and in consequence increase therapeutic compliance by its health promoting effect [24]. Small sample size, short duration of trial and gender-based sampling were among the weaknesses of the current assessment. Certainly, further methodical studies in future, with greater sample and longer duration, will prove more vividly various advantageous of this method in cardiac patients with psychiatric co-morbidity.

## Conclusion

Consistent with encouraging outcome of present study, cardiac rehabilitation can be recommended as a supplementary technique, in addition to other standard treatments, in management of ischemic heart patients with co-morbid depressive illness.

## References

1. Wayne Katon and Mark Sullivan. "Cardiovascular disorders". Comprehensive textbook of psychiatry; 6<sup>th</sup> edition (1995): 1491-1510.
2. KAPLAN and SADOCK. Psychological factors Affecting medical condition and psychosomatic medicine, synopsis of psychiatry, 9<sup>th</sup> edition (2003): 829-831.
3. KAPLAN and SADOCK. Late Adulthood (old age), synopsis of psychiatry, 9<sup>th</sup> edition (2003): 53.
4. Shoja shafti S. "Prevalence of Psychiatric Morbidities in Acute Coronary Heart Disease". *Cardiovascular Psychiatry and Neurology* 3.1 (2014):1- 5.
5. Haines AP, *et al.* "Phobic anxiety and ischemia heart disease". *The British Medical Journal* 295.6593 (1987): 297-299.
6. Kaprio J, *et al.* "Mortality after bereavement: a prospective study of 95,647 widowed persons". *The American Journal of Public Health* 77.3 (1987): 283-287.
7. Booth-Kewley S and Friedman HS. "Psychological predictors of heart disease: a quantitative review". *Psychological Bulletin* 101.3 (1987): 343-362.
8. Rosenman RH, *et al.* "Coronary heart disease in Western Collaborative Group Study: final follow-up experience of 8 ½ years". *The Journal of the American Medical Association* 233.8 (1975): 872-877.

---

**Citation:** Saeed Shoja Shafti MD and Masuod Ahmadi MD. "Effectiveness of Cardiac Rehabilitation on Co-Morbid Psychiatric Symptoms of Ischemic Heart Patients". *Current Opinions in Neurological Science* 3.1 (2019): 637-643.

9. Hecker MH., *et al.* "Coronary-prone behaviors in the Western Collaborative group study". *Psychosomatic Medicine* 50.2 (1988): 153-164.
10. Hearn MD., *et al.* "Hostility, coronary heart disease, and total mortality: a 33-year follow up study of university students". *Journal of Behavioral Medicine* 12.2 (1989): 105-121.
11. Hamilton M. "Rating Scale for Depression". *Journal of Neurology, Neurosurgery, and Psychiatry* 23 (1960): 56.
12. Schulman JK., *et al.* "Psychiatry and cardiovascular disease". *FOCUS* 3.2 (2005): 208-224.
13. Aben I., *et al.* "A comparative study into the one year cumulative incidence of depression after stroke and myocardial infarction". *Journal of Neurology Neurosurgery and Psychiatry* 74.5 (2003) 581- 585.
14. Farisa R., *et al.* "Clinical depression is common and significantly associated with reduced survival in patients with non-ischaemic heart failure". *European Journal of Heart Failure* 4.4 (2002): 541-551.
15. Moser DK and Dracup K. "Is anxiety early after myocardial infarction associated with subsequent ischemic and arrhythmic events?" *Psychosomatic Medicine* 58.5 (1996): 395-403.
16. Rudisch B and Nemeroff CB. "Epidemiology of comorbid coronary artery disease and depression". *Biological Psychiatry* 54.3 (2003): 227-240.
17. Rugulies R. "Depression as a predictor for coronary heart disease: a review and meta-analysis". *The American Journal of Preventive Medicine* 23.1 (2002): 51-61.
18. Gustad T., *et al.* "Symptoms of anxiety and depression and risk of acute myocardial infarction: the HUNT 2 study". *European Heart Journal* 35.3 (2014): 1394-1403.
19. Lavie CJ and Milani RV. "Adverse psychological and coronary risk profiles in young patients with coronary artery disease and benefits of formal cardiac rehabilitation". *Archives of Internal Medicine* 166 (2006): 1878-1883.
20. Milani RV., *et al.* "Impact of exercise training and depression on survival in heart failure due to coronary heart disease". *American Journal of Cardiology* 107.1 (2011): 64-68.
21. Linden W., *et al.* "Psychosocial interventions for patients with coronary artery disease: a meta-analysis". *Archives of Internal Medicine* 156 (1996): 745-752.
22. Johnston M and Foulkers J. "Impact on Patients and Partners of Inpatient and Extended Cardiac Counseling and Rehabilitation: A Controlled Trial". *Journal of Psychosomatic Medicine* 61.2 (1999): 225-233.
23. Braundwaid E: *Heart disease*, 6th edition 40 (2001):1411
24. Poshkov K., *et al.* *Clinical cardiac rehabilitation. A cardiologist guide*, Second edition, 1 (1999): 17.

**Submit your next manuscript to Scientia Ricerca Open Access and benefit from:**

- Prompt and fair double blinded peer review from experts
- Fast and efficient online submission
- Timely updates about your manuscript status
- Sharing Option: Social Networking Enabled
- Open access: articles available free online
- Global attainment for your research

Submit your manuscript at:

<https://scientiaricerca.com/submit-manuscript.php>